



FOREWORD

Shark, skate and ray research at the MBA and Cefas

Jim Ellis*, Nick Dulvy*, Carl O'Brien*, David Sims[†] and Emily Southall[†]

*Centre for Environment, Fisheries and Aquaculture Science (Cefas), Lowestoft Laboratory, Pakefield Road, Lowestoft, Suffolk, NR33 0HT, UK. [†]Marine Biological Association, The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK.

For many decades, scientists at the Marine Biological Association (MBA) in Plymouth and the Centre for Environment, Fisheries and Aquaculture Science (Cefas), formerly the Directorate of Fisheries Research, at Lowestoft have undertaken considerable research into the ecology and life histories of elasmobranch fish around the British Isles.

Based at Plymouth, E. Ford (1921) published one of the earliest accounts of the more common dogfish, including their population structure, reproductive biology and feeding habits. The first information on the breeding grounds of dogfish were also reported (Ford, 1921; Orton, 1926). At this time, R.S. Clark undertook several studies on the skates and rays occurring in the waters of the British Isles, focusing on the egg-capsules and early life history stages (Clark, 1922, 1927) and providing detailed accounts of their general biology and morphology (Clark, 1926). Subsequent studies on skates and rays were undertaken by G.A. Steven, who examined commercial catches and undertook studies on their feeding ecology, reproductive biology and movements (Steven, 1930, 1931, 1932, 1933, 1934, 1936, 1947). Around this time, much of the fisheries research conducted at Lowestoft was concerned with commercial teleosts, though C.F. Hickling undertook biological sampling of spurdog *Squalus acanthias* and other squaloids during a series of cruises collecting information on hake in the Celtic Sea (Hickling, 1930, 1963).

Commercial fisheries expanded greatly after the Second World War, including fisheries for spurdog and various skates and rays. During the 1960s and 1970s Lowestoft initiated a programme of elasmobranch research, which was led by M.J. Holden, who was one of the first scientists to highlight the susceptibility of elasmobranchs to over-exploitation (Holden, 1973, 1974, 1977). This period of research resulted in numerous publications regarding the age, growth, reproductive biology and feeding habits of a variety of skates and rays (Holden, 1963, 1972, 1975; Taylor & Holden, 1964; Holden et al., 1971; Holden & Vince, 1973; Holden & Tucker, 1974) and spurdog (Holden, 1960, 1964, 1965a,b, 1967a,b, 1968; Holden & Meadows, 1962, 1964; Tucker, 1985; Vince, 1991), and much of the biological knowledge gained from these studies remains invaluable today. Despite the increase in biological studies, effective management plans for elasmobranchs in European waters were not brought in, and subsequently Lowestoft scientists

provided the first documentation of the disappearance of common skate *Dipturus batis* from coastal waters of the United Kingdom (Brander, 1981).

Though the Lowestoft laboratory had a large research programme for skates, rays and coastal dogfish, less research had been focused on pelagic sharks. J.D. Stevens, working with Q. Bone at the MBA, began studies to improve our biological knowledge of blue shark *Prionace glauca* and other pelagic species in the early 1970s (e.g. Stevens, 1973, 1974, 1975, 1976, 1990). Our present knowledge of the movements, migrations and population structure of blue shark in the north-east Atlantic still relies heavily on this work. In more recent years, the MBA has undertaken much research into the movements and behaviour of several elasmobranchs, notably the basking shark *Cetorhinus maximus* and lesser-spotted dogfish *Scyliorhinus canicula* (Sims, 2003). Improved understanding of the free-ranging behaviour of basking sharks for instance, has contributed important new information relevant to national and international conservation efforts.

Since becoming an Executive Agency of the Department for Environment, Food and Rural Affairs (Defra) in 1997, Cefas has continued to undertake research on elasmobranchs, with Cefas scientists involved with the ICES Working Group on Elasmobranch Fish (WGEF), and the EC-funded project 'Development of Elasmobranch Assessments (DELASS)' (Heessen, 2003). There have also been MBA–Cefas research collaborations, most notably on Defra-funded satellite-tracking studies of basking sharks (Sims et al., 2003).

The biological characteristics of elasmobranchs, including a late age at maturity, slow growth rate, low fecundity and long gestation period, result in a low rate of population increase, thus making this group of fish particularly vulnerable to fishing pressure. Recent studies by Cefas and MBA scientists have documented long-term changes in the distribution and relative abundance of a variety of elasmobranchs in the waters around the British Isles, and related such changes to the intrinsic vulnerability and extrinsic mortality imposed by fishing (e.g. Dulvy et al., 2000; Rogers & Ellis, 2000; Genner et al., 2004; Ellis et al., 2005). Indeed several species of elasmobranch, such as angel shark *Squatina squatina* and white skate *Rostroraja alba*, have declined dramatically in the north-east Atlantic and are now rarely observed. Many

other species that are landed in mixed and targeted commercial fisheries have been over-fished, including spurdog (Hammond & Ellis, 2005) and various skates, and appropriate management measures are required to ensure the sustainability of these stocks. It should also be noted that many elasmobranchs are important target species for recreational fisheries and, from a socio-economic aspect, some species may be more valuable to these fisheries than to the commercial fleet.

Elasmobranch fish possess several biological characteristics that make them of fundamental scientific interest. For example, they have advanced sensory modalities, relatively large and structurally complex brains, the capability for complex behaviours such as social systems and sexual segregation, internal fertilization and diverse reproductive strategies ranging from egg-laying to live-bearing. Scientific research is shedding light on these and many other aspects of elasmobranch biology. An improved biological knowledge of these species is a fundamental requirement for the management of elasmobranch stocks, especially with regards to greater understanding of their movements, reproductive biology and population structure.

There are 23 papers on elasmobranchs in this special issue of the *Journal of the Marine Biological Association*, discussing a range of elasmobranch species and various aspects of their biology, including migrations, reproductive biology, population structure and diet. Migrations and movements are discussed for several species, including thornback ray *Raja clavata* in the outer Thames Estuary (Hunter et al.), greater-spotted dogfish *Scyliorhinus stellaris* (Sims et al.), basking shark *Cetorhinus maximus* (Southall et al.) and blue shark *Prionace glauca* (Queiroz et al.). Aspects of the reproductive biology are discussed for narrownose smooth hound *Mustelus schmitti* in the south-east Atlantic (Oddone et al.), black-mouth catshark *Galeus melastomus* in Portuguese waters (Costa et al.), various skates and rays (Koop), spurdog *Squalus acanthias* in the eastern Mediterranean (Chatzisprou & Megalofonou), Port Jackson shark *Heterodontus portusjacksoni* (Jones et al.), lantern sharks *Etmopterus* spp. (Coelho & Erzini) and a possible new species of gulper shark *Centrophorus* cf. *uyato* from the Cayman Trench (McLaughlin & Morrissey). Age, growth and other aspects of the population structure have been discussed for *R. clavata* in the Irish Sea (Whittamore & McCarthy), *Psammobatis extenta* off Brazil (Martins et al.), and sandbar shark *Carcharhinus plumbeus* (Torres et al.). The feeding habits of various elasmobranchs are discussed, such as white shark *Carcharodon carcharias* (Martin et al.) and the electric ray *Discopyge tshudii* (Arrighetti et al.).

Freshwater and estuarine elasmobranchs are often regarded as of particular conservation concern, as reviewed by Martin. Both fisheries and habitat degradation, can affect populations of these species, especially since their distributions are relatively small. The low fecundity of many deep-water sharks (e.g. McLaughlin & Morrissey) means that many of these species may need very conservative management if they are able to support fisheries, especially since our current knowledge of the stock structure of deep-water elasmobranchs is so poor. Fisheries management for some elasmobranchs is slowly improving, but is hampered by a lack of species-specific landings data, and uncertainty in both stock identity and

population structure, and so the assessment methods widely applied to commercial teleosts cannot yet be applied to most elasmobranchs. Demersal elasmobranchs may be sampled in trawl surveys (Ellis et al.), though not all species and life history stages are sampled effectively, and catch rates can be highly variable. Given that many demersal elasmobranchs, which include many species of conservation concern, are taken in mixed fisheries, improved estimates of survivorship are required if quotas or size restrictions are to be effective management measures. Rodríguez-Cabello et al. have shown that *Scyliorhinus canicula* discarded from trawlers have a high survivorship, though estimates of discard mortality are lacking for most other species and gears. In addition to fisheries impacts on elasmobranch stocks, the recent expansion of wind farms in marine environments and the potential impacts of such activities on fish and fisheries has attracted much interest in recent years, as reviewed by Gill & Kimber.

An improved biological knowledge of elasmobranchs is required to facilitate appropriate management of their stocks, so that the biodiversity of this group of fish can be maintained, and that commercial and recreational fisheries for elasmobranchs can be managed sustainably. The MBA and Cefas are pleased to produce this special issue on elasmobranch fish, with many of these papers arising from the 8th European Elasmobranch Association Conference, held from 21–24 October 2004 at the Zoological Society of London, and organized by the Shark Trust (www.sharktrust.org) and Zoological Society of London (<http://www.zsl.org/>). To complete this special issue, various papers on teleost fish have also been included. All the papers have been subjected to *JMBA* review.

REFERENCES

- Brander, K., 1981. Disappearance of common skate *Raia batis* from Irish Sea. *Nature, London*, **290**, 48–49.
- Clark, R.S., 1922. Rays and skates (*Raiae*). No. I: egg capsules and young. *Journal of the Marine Biological Association of the United Kingdom*, **12**, 577–643.
- Clark, R.S., 1926. Rays and skates. A revision of the European species. *Scientific Investigations, Fishery Board for Scotland*, **1**, 1–66.
- Clark, R.S., 1927. Rays and skates. No. 2: description of embryos. *Journal of the Marine Biological Association of the United Kingdom*, **14**, 661–683.
- Dulvy, N.K., Metcalfe, J.D., Glanville, J., Pawson, M.G. & Reynolds, J.D., 2000. Fishery stability, local extinctions, and shifts in community structure in skates. *Conservation Biology*, **14**, 283–293.
- Ellis, J.R., Cruz-Martinez, A., Rackham, B.D. & Rogers, S.I., 2005. The distribution of chondrichthyan fishes around the British Isles and implications for conservation. *Journal of Northwest Atlantic Fishery Science*, **35**, 195–213.
- Ford, E., 1921. A contribution to our knowledge of the life-histories of the dogfishes landed at Plymouth. *Journal of the Marine Biological Association of the United Kingdom*, **12**, 468–505.
- Genner, M.J., Sims, D.W., Wearmouth, V.J., Southall, E.J., Southward, A.J., Henderson, P.A. & Hawkins, S.J., 2004. Regional climate warming drives long-term community changes of British marine fish. *Proceedings of the Royal Society of London B*, **271**, 655–661.
- Hammond, T.R. & Ellis, J.R., 2005. Bayesian assessment of North-east Atlantic spurdog using a stock production model, with prior for intrinsic population growth rate set by demographic methods. *Journal of Northwest Atlantic Fishery Science*, **35**, 299–308.

- Heessen, H.J.L., ed., 2003. Development of Elasmobranch Assessments—DELIASS: DG Fish Study Contract 99/055.
- Hickling, C.F., 1930. A contribution towards the life history of the spurdog. *Journal of the Marine Biological Association of the United Kingdom*, **16**, 529–576.
- Hickling, C.F., 1963. On the small deep-sea shark *Etmopterus spinax* L. and its cirripede parasite *Anelasma squalicola* (Loven). *Journal of the Linnean Society (Zoology)*, **45**, 17–24.
- Holden, M.J., 1960. English spurdogfish tagging. *Annales Biologiques, Copenhagen*, **17**, 107–108.
- Holden, M.J., 1963. The species composition of skates and rays landed at Fleetwood and Milford Haven. *ICES CM 1963 Near Northern Seas Committee*, no. **57**, 3 pp.
- Holden, M.J., 1964. English taggings of picked dogfish. *Annales Biologiques, Copenhagen*, **21**, 172–174.
- Holden, M.J., 1965a. The food of the spurdog, *Squalus acanthias* (L.). *Journal du Conseil*, **30**, 255–266.
- Holden, M.J., 1965b. The stocks of spurdogs (*Squalus acanthias* L.) in British waters, and their migrations. *Fishery Investigations, London*, Series 2, **24**(4), 20 pp.
- Holden, M.J., 1967a. Spurdogs. Laboratory Leaflet (New Series), MAFF Fisheries Laboratory, Lowestoft, 23 pp.
- Holden, M.J., 1967b. Transatlantic movement of a tagged spurdogfish. *Nature, London*, **214**, 1140–1141.
- Holden, M.J., 1968. The rational exploitation of the Scottish–Norwegian stock of spurdogs (*Squalus acanthias* L.). *Fishery Investigations, London*, Series 2, **25**(8), 27 pp.
- Holden, M.J., 1972. The growth rates of *Raja brachyura*, *R. clavata* and *R. montagui* as determined from tagging data. *Journal du Conseil*, **34**, 161–168.
- Holden, M.J., 1973. Are long-term sustainable fisheries for elasmobranchs possible? *Rapports et Procès-verbaux des Réunions, Conseil International pour l'Exploration de la Mer*, **164**, 360–367.
- Holden, M.J., 1974. Problems in the rational exploitation of elasmobranch populations and some suggested solutions. In *Sea fisheries research* (ed. F.R. Harden Jones), pp. 117–137. Elek: London.
- Holden, M.J., 1975. The fecundity of *Raja clavata* in British waters. *Journal du Conseil*, **36**, 110–118.
- Holden, M.J., 1977. Elasmobranchs. In *Fish population dynamics* (ed. G.A. Gulland), pp. 187–215. London: John Wiley & Sons.
- Holden, M.J. & Meadows, P.S., 1962. The structure of the spine of the spurdog fish (*Squalus acanthias* L.) and its use for age determination. *Journal of the Marine Biological Association of the United Kingdom*, **42**, 179–197.
- Holden, M.J. & Meadows, P.S., 1964. The fecundity of the spurdog (*Squalus acanthias* L.). *Journal du Conseil*, **28**, 418–424.
- Holden, M.J., Rout, D.W. & Humphreys, C.N., 1971. The rate of egg laying by three species of ray. *Journal du Conseil*, **33**, 335–339.
- Holden, M.J. & Tucker, R.N., 1974. The food of *Raja clavata*, *R. montagui*, *R. naevus* and *R. brachyura* in British waters. *Journal du Conseil*, **35**, 189–193.
- Holden, M.J. & Vince, M.R., 1973. Age validation studies on the centra of *Raja clavata* using tetracycline. *Journal du Conseil*, **35**, 13–17.
- Orton, J.H., 1926. A breeding ground of the nursehound (*Scyliorhinus stellaris*) in the Fal estuary. *Nature, London*, **118**, 732.
- Rogers, S.I. & Ellis, J.R., 2000. Changes in the demersal fish assemblages of British coastal waters during the 20th century. *ICES Journal of Marine Science*, **57**, 866–881.
- Sims, D.W., 2003. Tractable models for testing theories about natural strategies: foraging behaviour and habitat selection of free-ranging sharks. *Journal of Fish Biology*, **63**, Supplement A, 53–73.
- Sims, D.W., Southall, E.J., Richardson, A.J., Reid, P.C. & Metcalfe, J.D., 2003. Seasonal movements and behaviour of basking sharks from archival tagging: no evidence of winter hibernation. *Marine Ecology Progress Series*, **248**, 187–196.
- Steven, G.A., 1930. Bottom fauna and the food of fishes. *Journal of the Marine Biological Association of the United Kingdom*, **16**, 677–706.
- Steven, G.A., 1931. Rays and skates of Devon and Cornwall. Methods of rapid identification on the fishmarket. *Journal of the Marine Biological Association of the United Kingdom*, **17**, 367–377.
- Steven, G.A., 1932. Rays and skates of Devon and Cornwall II. A study of the fishery; with notes on the occurrence, migrations and habits of the species. *Journal of the Marine Biological Association of the United Kingdom*, **18**, 1–33.
- Steven, G.A., 1933. Rays and skates of Devon and Cornwall III. The proportions of the sexes in nature and in commercial landings and their significance to the fishery. *Journal of the Marine Biological Association of the United Kingdom*, **18**, 611–625.
- Steven, G.A., 1934. Observations on the growth of the claspers and cloaca in *Raja clavata* Linnaeus. *Journal of the Marine Biological Association of the United Kingdom*, **19**, 887–899.
- Steven, G.A., 1936. Migrations and growth of the thornback ray (*Raja clavata* L.). *Journal of the Marine Biological Association of the United Kingdom*, **20**, 605–614.
- Steven, G.A., 1947. The British Rajidae. *Science Progress*, **35**, 220–236.
- Stevens, J.D., 1973. Stomach contents of the blue shark (*Prionace glauca* L.) off south-west England. *Journal of the Marine Biological Association of the United Kingdom*, **53**, 357–361.
- Stevens, J.D., 1974. The occurrence and significance of tooth cuts on the blue shark (*Prionace glauca* L.) from British waters. *Journal of the Marine Biological Association of the United Kingdom*, **54**, 373–378.
- Stevens, J.D., 1975. Vertebral rings as a means of age determination in the blue shark (*Prionace glauca*). *Journal of the Marine Biological Association of the United Kingdom*, **55**, 657–665.
- Stevens, J.D., 1976. First results of shark tagging in the north-east Atlantic, 1972–1975. *Journal of the Marine Biological Association of the United Kingdom*, **56**, 929–937.
- Stevens, J.D., 1990. Further results from a tagging study of pelagic sharks in the north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom*, **70**, 707–720.
- Taylor, A.J. & Holden, M.J., 1964. The preparation and use of vertebrae for age determination in rays. *ICES CM 1964*, no. 145, 5 pp.
- Tucker, R., 1985. Age validation studies on the spines of the spurdog (*Squalus acanthias*) using tetracycline. *Journal of the Marine Biological Association of the United Kingdom*, **65**, 641–651.
- Vince, M.R., 1991. Stock identity in spurdog (*Squalus acanthias* L.) around the British Isles. *Fisheries Research*, **12**, 341–354.